

## References

1. Gore AC, Chappell VA, Fenton SE, et al. EDC-2: The Endocrine Society's Second Scientific Statement on Endocrine-Disrupting Chemicals. *Endocr Rev.* 2015 Dec;36(6):E1-E150.
2. Moos RK, Angerer J, Dierkes G, Brüning T, Koch HM. Metabolism and elimination of methyl, iso- and n-butyl paraben in human urine after single oral dosage. *Arch Toxicol.* 2016 Nov;90(11):2699–709.
3. Völkel W, Colnot T, Csanády GA, Filser JG, Dekant W. Metabolism and kinetics of bisphenol a in humans at low doses following oral administration. *Chem Res Toxicol.* 2002 Oct;15(10):1281–7.
4. Anderson WA, Castle L, Scotter MJ, Massey RC, Springall C. A biomarker approach to measuring human dietary exposure to certain phthalate diesters. *Food Addit Contam.* 2001 Dec;18(12):1068–74.
5. Bellou V, Belbasis L, Tzoulaki I, Evangelou E. Risk factors for type 2 diabetes mellitus: An exposure-wide umbrella review of meta-analyses. *PLoS One.* 2018 Mar 20;13(3):e0194127.
6. Fukata H, Miyagawa H, Yamazaki N, Mori C. Comparison of Elisa- and LC-MS-Based Methodologies for the Exposure Assessment of Bisphenol A. *Toxicol Mech Methods.* 2006;16(8):427–30.
7. Koch HM, Gonzalez-Reche LM, Angerer J. On-line clean-up by multidimensional liquid chromatography-electrospray ionization tandem mass spectrometry for high throughput quantification of primary and secondary phthalate metabolites in human urine. *J Chromatogr B Analyt Technol Biomed Life Sci.* 2003;784(1):169–82.
8. Silva MJ, Samandar E, Preau JL, Reidy JA, Needham LL, Calafat AM. Quantification of 22 phthalate metabolites in human urine. *J Chromatogr B Analyt Technol Biomed Life Sci.* 2007;860(1):106–12.
9. Kato K, Silva MJ, Needham LL, Calafat AM. Determination of 16 phthalate metabolites in urine using automated sample preparation and on-line preconcentration/highperformance liquid chromatography/tandem mass spectrometry. *Anal Chem.* 2005;77(9):2985–91.
10. Heffernan AL, Thompson K, Eaglesham G, Vijayasarathy S, Mueller JF, Sly PD, et al. Rapid, automated online SPE-LC-QTRAP-MS/MS method for the simultaneous analysis of 14 phthalate metabolites and 5 bisphenol analogues in human urine. *Talanta.* 2016;151:224–33.
11. Ye X, Kuklennyik Z, Needham LL, Calafat AM. Automated on-line column-switching HPLC-MS/MS method with peak focusing for the determination of nine environmental phenols in urine. *Anal Chem.* 2005;77(16):5407–13.
12. Zhou X, Kramer JP, Calafat AM, Ye X. Automated on-line column-switching high performance liquid chromatography isotope dilution tandem mass spectrometry method for the quantification of bisphenol A, bisphenol F, bisphenol S, and 11 other phenols in urine. *J Chromatogr B Analyt Technol Biomed Life Sci.* 2014;944:152–6.

13. Schindler BK, Esteban M, Koch HM, Castano A, Koslitz S, Cañas A, et al. The European COPHES/DEMOCOPHES project: towards transnational comparability and reliability of human biomonitoring results. *Int J Hyg Environ Health*. 2014 Jul;217(6):653–61.
14. Moos RK, Angerer J, Dierkes G, Brüning T, Koch HM. Metabolism and elimination of methyl, iso- and n-butyl paraben in human urine after single oral dosage. *Arch Toxicol*. 2016 Nov;90(11):2699–709.
15. Völkel W, Colnot T, Csanády GA, Filser JG, Dekant W. Metabolism and kinetics of bisphenol a in humans at low doses following oral administration. *Chem Res Toxicol*. 2002 Oct;15(10):1281–7.
16. Anderson WA, Castle L, Scotter MJ, Massey RC, Springall C. A biomarker approach to measuring human dietary exposure to certain phthalate diesters. *Food Addit Contam*. 2001 Dec;18(12):1068–74.
17. Christensen KL, Lorber M, Koch HM, Kolossa-Gehring M, Morgan MK. Population variability of phthalate metabolites and bisphenol A concentrations in spot urine samples versus 24- or 48-h collections. *J Expo Sci Environ Epidemiol*. 2012;22(6):632–40.
18. Johns LE, Cooper GS, Galizia A, Meeker JD. Exposure assessment issues in epidemiology studies of phthalates. *Environ Int*. 2015 Dec;85:27–39.
19. Preau JL Jr, Wong L-Y, Silva MJ, Needham LL, Calafat AM. Variability over 1 week in the urinary concentrations of metabolites of diethyl phthalate and di(2-ethylhexyl) phthalate among eight adults: an observational study. *Environ Health Perspect*. 2010 Dec;118(12):1748–54.
20. LaKind JS, Naiman DQ. Temporal trends in bisphenol A exposure in the United States from 2003–2012 and factors associated with BPA exposure: Spot samples and urine dilution complicate data interpretation. *Environ Res*. 2015;142:84–95.
21. Koch HM, Rüter M, Schütze A, Conrad A, Pälme C, Apel P, et al. Phthalate metabolites in 24-h urine samples of the German Environmental Specimen Bank (ESB) from 1988 to 2015 and a comparison with US NHANES data from 1999 to 2012. *Int J Hyg Environ Health*. 2017 Mar;220(2 Pt A):130–41.
22. LaKind JS, Pollock T, Naiman DQ, Kim S, Nagasawa A, Clarke J. Factors affecting interpretation of national biomonitoring data from multiple countries: BPA as a case study. *Environ Res*. 2019 Jun;173:318–29.
23. Frederiksen H, Jensen TK, Jørgensen N, Kyhl HB, Husby S, Skakkebaek NE, et al. Human urinary excretion of non-persistent environmental chemicals: an overview of Danish data collected between 2006 and 2012. *Reproduction*. 2014;147(4):555–65.
24. Frederiksen H, Nielsen O, Koch HM, Skakkebaek NE, Juul A, Jørgensen N, et al. Changes in urinary excretion of phthalates, phthalate substitutes, bisphenols and other polychlorinated and phenolic substances in young Danish men; 2009–2017. *Int J Hyg Environ Health*. 2020 Jan;223(1):93–105.

25. Ye X, Pierik FH, Hauser R, Duty S, Angerer J, Park MM, et al. Urinary metabolite concentrations of organophosphorous pesticides, bisphenol A, and phthalates among pregnant women in Rotterdam, the Netherlands: the Generation R study. *Environ Res.* 2008;108(2):260–7.
26. Philips EM, Jaddoe VWV, Asimakopoulos AG, Kannan K, Steegers EAP, Santos S, et al. Bisphenol and phthalate concentrations and its determinants among pregnant women in a population-based cohort in the Netherlands, 2004-5. *Environ Res.* 2018;161:562–72.
27. Philips EM, Jaddoe VWV, Deierlein A, Asimakopoulos AG, Kannan K, Steegers EAP, et al. Exposures to phthalates and bisphenols in pregnancy and postpartum weight gain in a population-based longitudinal birth cohort. *Environ Int.* 2020 Jul 31;144:106002.
28. COMMISSION REGULATION (EU) No 358/2014 [Internet]. Official Journal of the European Union. 2014 [cited 2020 Jun 18]. Available from: <https://eurlex.europa.eu/legal-content/EN/TXT/HTML/?uri=CELEX:32014R0358&from=EN>
29. EFSA Panel on Food Contact Materials, Enzymes, Flavourings And Processing. Scientific Opinion on the risks to public health related to the presence of bisphenol A (BPA) in foodstuffs. *EFSA Journal.* 2015 Jan 1;13(1):3978.
30. Ye X, Wong L-Y, Kramer J, Zhou X, Jia T, Calafat AM. Urinary Concentrations of Bisphenol A and Three Other Bisphenols in Convenience Samples of U.S. Adults during 2000–2014. *Environ Sci Technol.* 2015 Oct 6;49(19):11834–9.
31. COMMISSION REGULATION (EU) 2016/2235 [Internet]. Available from: <https://eurlex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32016R2235&from=EN>
32. COMMISSION REGULATION (EU) 2018/2005 [Internet]. Available from: <https://eurlex.europa.eu/legal-content/EN/TXT/PDF/?uri=CELEX:32018R2005&from=EN>
33. Directive 2011/65/EU [Internet]. Available from: <https://eur-lex.europa.eu/legalcontent/EN/TXT/PDF/?uri=CELEX:32015L0863&from=FR>
34. Rochester JR, Bolden AL. Bisphenol S and F: A Systematic Review and Comparison of the Hormonal Activity of Bisphenol A Substitutes. *Environ Health Perspect.* 2015;123(7):643–50.
35. Aubert N, Ameller T, Legrand J-J. Systemic exposure to parabens: pharmacokinetics, tissue distribution, excretion balance and plasma metabolites of [14C]-methyl-, propyl and butylparaben in rats after oral, topical or subcutaneous administration. *Food Chem Toxicol.* 2012 Mar;50(3-4):445–54.
36. Janjua NR, Frederiksen H, Skakkebaek NE, Wulf HC, Andersson A-M. Urinary excretion of phthalates and paraben after repeated whole-body topical application in humans. *Int J Androl.* 2008 Apr;31(2):118–30.
37. Bhagwat VM, Ramachandran BV. Malathion A and B esterases of mouse liver-I. *Biochem Pharmacol.* 1975 Sep 15;24(18):1713–7.
38. Christensen KLY, Lorber M, Koslitz S, Brüning T, Koch HM. The contribution of diet to

total bisphenol A body burden in humans: results of a 48 hour fasting study. *Environ Int.* 2012 Dec 1;50:7–14.

39. Rudel RA, Gray JM, Engel CL, Rawsthorne TW, Dodson RE, Ackerman JM, et al. Food packaging and bisphenol A and bis(2-ethylhexyl) phthalate exposure: findings from a dietary intervention. *Environ Health Perspect.* 2011 Jul;119(7):914–20.

40. Koch HM, Lorber M, Christensen KLY, Pälme C, Koslitz S, Brüning T. Identifying sources of phthalate exposure with human biomonitoring: results of a 48h fasting study with urine collection and personal activity patterns. *Int J Hyg Environ Health.* 2013 Nov;216(6):672–81.

41. Serrano SE, Braun J, Trasande L, Dills R, Sathyanarayana S. Phthalates and diet: a review of the food monitoring and epidemiology data. *Environ Health.* 2014;13(1):43.

42. National Research Council. *Phthalates and Cumulative Risk Assessment.* 2008.

43. Dirtu AC, Geens T, Dirinck E, Malarvannan G, Neels H, Van Gaal L, et al. Phthalate metabolites in obese individuals undergoing weight loss: Urinary levels and estimation of the phthalates daily intake. *Environ Int.* 2013 Sep;59:344–53.

44. Colacino JA, Harris TR, Schecter A. Dietary intake is associated with phthalate body burden in a nationally representative sample. *Environ Health Perspect.* 2010 Jul;118(7):998–1003.

45. Trasande L, Sathyanarayana S, Jo Messito M, S Gross R, Attina TM, Mendelsohn AL. Phthalates and the diets of U.S. children and adolescents. *Environ Res.* 2013 Oct;126:84–90.

46. Lind PM, Roos V, Rönn M, Johansson L, Ahlström H, Kullberg J, et al. Serum concentrations of phthalate metabolites are related to abdominal fat distribution two years later in elderly women. *Environ Health.* 2012 Apr 2;11:21.

47. LaKind JS, Idri F, Naiman DQ, Verner M-A. Biomonitoring and Nonpersistent Chemicals-Understanding and Addressing Variability and Exposure Misclassification. *Curr Environ Health Rep.* 2019 Mar;6(1):16–21.

48. Starling AP, Engel LS, Calafat AM, Koutros S, Satagopan JM, Yang G, et al. Predictors and long-term reproducibility of urinary phthalate metabolites in middle-aged men and women living in urban Shanghai. *Environ Int.* 2015 Nov;84:94–106.

49. Townsend MK, Franke AA, Li X, Hu FB, Eliassen AH. Within-person reproducibility of urinary bisphenol A and phthalate metabolites over a 1 to 3 year period among women in the Nurses' Health Studies: a prospective cohort study. *Environ Health.* 2013 Sep 13;12(1):80.

50. Sun Q, Cornelis MC, Townsend MK, Tobias DK, Eliassen AH, Franke AA, et al. Association of urinary concentrations of bisphenol A and phthalate metabolites with risk of type 2 diabetes: a prospective investigation in the Nurses' Health Study (NHS) and NHSII cohorts. *Environ Health Perspect.* 2014 Jun;122(6):616–23.

51. Shu X, Tang S, Peng C, Gao R, Yang S, Luo T, et al. Bisphenol A is not associated with

a 5-year incidence of type 2 diabetes: a prospective nested case-control study. *Acta Diabetol.* 2018 Apr;55(4):369–75.

52. Salamanca-Fernández E, Iribarne-Durán LM, Rodríguez-Barranco M, Vela-Soria F, Olea N, Sánchez-Pérez MJ, et al. Historical exposure to non-persistent environmental pollutants and risk of type 2 diabetes in a Spanish sub-cohort from the European Prospective Investigation into Cancer and Nutrition study. *Environ Res.* 2020 Jun;185:109383.

53. Rancière F, Botton J, Slama R, Lacroix MZ, Debrauwer L, Charles MA, et al. Exposure to Bisphenol A and Bisphenol S and Incident Type 2 Diabetes: A Case-Cohort Study in the French Cohort D.E.S.I.R. *Environ Health Perspect.* 2019 Oct;127(10):107013.

54. Kolatorova L, Sramkova M, Vitku J, Vcelak J, Lischkova O, Starka L, et al. Parabens and their relation to obesity. *Physiol Res.* 2018 Nov 28;67(Suppl 3):S465–72.

55. Kang H-S, Kyung M-S, Ko A, Park J-H, Hwang M-S, Kwon J-E, et al. Urinary concentrations of parabens and their association with demographic factors: A population-based cross-sectional study. *Environ Res.* 2016 Apr;146:245–51.

56. Quirós-Alcalá L, Buckley JP, Boyle M. Parabens and measures of adiposity among adults and children from the U.S. general population: NHANES 2007-2014. *Int J Hyg Environ Health.* 2018 May;221(4):652–60.

57. Trasande L, Attina TM, Blustein J. Association between urinary bisphenol A concentration and obesity prevalence in children and adolescents. *JAMA.* 2012 Sep 19;308(11):1113–21.

58. Liu B, Lehmler H-J, Sun Y, Xu G, Liu Y, Zong G, et al. Bisphenol A substitutes and obesity in US adults: analysis of a population-based, cross-sectional study. *Lancet Planet Health.* 2017 Jun;1(3):e114–22.

59. Mouneimne Y, Nasrallah M, Khoueiry-Zgheib N, Nasreddine L, Nakhoul N, Ismail H, et al. Bisphenol A urinary level, its correlates, and association with cardiometabolic risks in Lebanese urban adults. *Environ Monit Assess.* 2017 Oct 23;189(10):517.

60. James-Todd TM, Huang T, Seely EW, Saxena AR. The association between phthalates and metabolic syndrome: the National Health and Nutrition Examination Survey 2001-2010. *Environ Health.* 2016;15:52.

61. Hatch EE, Nelson JW, Qureshi MM, Weinberg J, Moore LL, Singer M, et al. Association of urinary phthalate metabolite concentrations with body mass index and waist circumference: a cross-sectional study of NHANES data, 1999-2002. *Environ Health.* 2008 Jun 3;7:27.

62. Stahlhut RW, van Wijngaarden E, Dye TD, Cook S, Swan SH. Concentrations of urinary phthalate metabolites are associated with increased waist circumference and insulin resistance in adult U.S. males. *Environ Health Perspect.* 2007 Jun;115(6):876–82.

63. Wang B, Li M, Zhao Z, Lu J, Chen Y, Xu Y, et al. Urinary bisphenol A concentration and glucose homeostasis in non-diabetic adults: a repeated-measures, longitudinal study. *Diabetologia.* 2019 Sep;62(9):1591–600.

64. Williams G, Harrold JA, Cutler DJ. The hypothalamus and the regulation of energy homeostasis: lifting the lid on a black box. *Proc Nutr Soc.* 2000 Aug;59(3):385–96.
65. Jandacek RJ, Anderson N, Liu M, Zheng S, Yang Q, Tso P. Effects of yo-yo diet, caloric restriction, and olestra on tissue distribution of hexachlorobenzene. *Am J Physiol Gastrointest Liver Physiol.* 2005 Feb;288(2):G292–9.
66. Imbeault P, Tremblay A, Simoneau J-A, Joanisse DR. Weight loss-induced rise in plasma pollutant is associated with reduced skeletal muscle oxidative capacity. *American Journal of Physiology-Endocrinology and Metabolism.* 2002 Mar 1;282(3):E574–9.
67. Malarvannan G, Van Hoorenbeeck K, Deguchtanaere A, Verhulst SL, Dirinck E, Van Gaal L, et al. Dynamics of persistent organic pollutants in obese adolescents during weight loss. *Environ Int.* 2018 Jan;110:80–7.
68. Artacho-Cordón F, Arrebola JP, Nielsen O, Hernández P, Skakkebaek NE, Fernández MF, et al. Assumed non-persistent environmental chemicals in human adipose tissue; matrix stability and correlation with levels measured in urine and serum. *Environ Res.* 2017;156:120–7.
69. Wang L, Asimakopoulos AG, Kannan K. Accumulation of 19 environmental phenolic and xenobiotic heterocyclic aromatic compounds in human adipose tissue. *Environ Int.* 2015;78:45–50.
70. Mes J, Coffin DE, Campbell DS. Di-n-butyl-and di-2-ethylhexyl phthalate in human adipose tissue. *Bull Environ Contam Toxicol.* 1974 Dec;12(6):721–5.
71. Mathieu-Denoncourt J, Wallace SJ, de Solla SR, Langlois VS. Influence of Lipophilicity on the Toxicity of Bisphenol A and Phthalates to Aquatic Organisms. *Bull Environ Contam Toxicol.* 2016 Jul;97(1):4–10.
72. Abbasi A, Peelen LM, Corpeleijn E, van der Schouw YT, Stolk RP, Spijkerman AMW, et al. Prediction models for risk of developing type 2 diabetes: systematic literature search and independent external validation study. *BMJ.* 2012 Sep 18;345:e5900.
73. Noble D, Mathur R, Dent T, Meads C, Greenhalgh T. Risk models and scores for type 2 diabetes: systematic review. *BMJ.* 2011 Nov 28;343:d7163.